



<http://jams.la.coocan.jp>

THE OMAN DRILLING PROJECT: A MILESTONE IN THE "MOHOLE TO MANTLE" (M2M) PROJECT

Japanese scientists and Japan's deep-sea scientific drilling vessel *Chikyū* are deeply involved in the Oman Drilling Project (Oman DP; <http://www.omandrilling.ac.uk/>), which is itself part of the International Continental Scientific Drilling Program (ICDP). From December 2016 to March 2017, Phase I drilling of the southern Oman ophiolite (Fig. 1) resulted in a total of 1,500 m of hard-rock cores being recovered. Measurements of the geophysical properties and geological description of the cores will be carried out on board the *Chikyū* in the summer of 2017 by shipboard scientists who applied to be involved from all over the world. The project has been nicknamed ChikyūOman. These cores have been sent to Japan for further study.




FIGURE 1 Simplified geologic map showing the locations of the Oman ophiolite and Oman Drilling Project drilling sites.

MODIFIED FROM NICOLAS ET AL. (2000).

The crust–mantle boundary, called the Moho Transition Zone (MTZ), in this region is thought to represent an active reaction zone where melts from a mid-ocean ridge basalt (MORB) accumulated and reacted with peridotites to form dunites (e.g. Boudier and Nicolas 1995; Kelemen et al. 1997; Korenaga and Kelemen 1997; Koga et al. 2001; Akizawa et al. 2012; Nicolle et al. 2016; Rospabé et al. 2017). Drilling into the MTZ is planned as Phase II of the Oman DP, and should take place from November 2017 to March 2018 (Fig. 1). The relationship between the MTZ and the major seismic-reflection boundary that is the Mohorovičić (Moho) discontinuity is still open to debate. The cores drilled through the Oman MTZ, and the geophysical logging of the boreholes, will provide a unique opportunity to evaluate and clarify the relationship between the Moho and the MTZ. This data will be essential for any future Mohole projects – drilling from the crust down to the Moho – one of which may be part of a future International Ocean Discovery Program (IODP) drilling project. The ultimate goal of the Oman DP is to help carry out the IODP's "Mohole to Mantle" (M2M) project, which aims to drill through a complete section of oceanic crust in the Pacific Ocean and into the underlying mantle (Kelemen et al. 2013). In this regard, the Oman DP and ChikyūOman can together be regarded as a milestone in the progress towards the M2M project.

Eiichi Takazawa
Niigata University / JAMSTEC

REFERENCES

- Akizawa N, Arai S, Tamura A (2012) Behavior of MORB magmas at uppermost mantle beneath a fast-spreading axis: an example from Wadi Fizh of the northern Oman ophiolite. *Contributions to Mineralogy and Petrology* 164: 601-625
- Boudier F, Nicolas A (1995) Nature of the Moho transition zone in the Oman ophiolite. *Journal of Petrology* 36: 777-796
- Kelemen PB, Koga K, Shimizu N (1997) Geochemistry of gabbro sills in the crust–mantle transition zone of the Oman ophiolite: implications for the origin of the oceanic lower crust. *Earth and Planetary Science Letters* 146: 475-488
- Kelemen P and 11 coauthors (2013) Scientific drilling and related research in the Samail ophiolite, Sultanate of Oman. *Scientific Drilling* 15: 64-71
- Koga KT, Kelemen PB, Shimizu N (2001) Petrogenesis of the crust–mantle transition zone and the origin of lower crustal wehrlite in the Oman ophiolite. *Geochemistry, Geophysics, Geosystems* 2: doi: 10.1029/2000GC000132
- Korenaga J, Kelemen PB (1997) Origin of gabbro sills in the Moho transition zone of the Oman ophiolite: implications for magma transport in the oceanic lower crust. *Journal of Geophysical Research* 102: 27729-27749
- Nicolas A, Boudier F, Ildefonse B, Ball E (2000) Accretion of Oman and United Arab Emirates ophiolite – discussion of a new structural map. *Marine Geophysical Researches* 21: 147-180
- Nicolle M, Jousset D, Reisberg L, Bosch D, Stephant A (2016) Major and trace element and Sr and Nd isotopic results from mantle diapirs in the Oman ophiolite: implications for off-axis magmatic processes. *Earth and Planetary Science Letters* 437: 138-149
- Rospabé M, Ceuleneer G, Benoit M, Abily B, Pinet P (2017) Origin of the dunitic mantle–crust transition zone in the Oman ophiolite: the interplay between percolating magmas and high-temperature hydrous fluids. *Geology* 45: 471-474

JOURNAL OF MINERALOGICAL AND PETROLOGICAL SCIENCES

Vol. 112, No. 3, June 2017

Original Articles

Abuite, $\text{CaAl}_2(\text{PO}_4)_2\text{F}_2$, a new mineral from the Hinomaru-Nago mine, Yamaguchi Prefecture, Japan – Satomi ENJU and Seiichiro UEHARA

Precipitates within olivine phenocrysts in oxidized andesitic scoria from Kasayama volcano, Hagi, Japan – Terumi EJIMA, Mari YONEDA, Masahide AKASAKA, Hiroaki OHFUJI, Yoshiaki KON, Mariko NAGASHIMA and Yoshihiro NAKAMUTA

Letters

Sintering nanocrystalline diopside from pulverized diopside crystals – Yumiko TSUBOKAWA and Masahiro ISHIKAWA

Carbonic fluid inclusions in a garnet–pyroxene granulite from Austhovde in the Lützow–Holm Complex, East Antarctica: Implications for a decompressional P – T path – Kazuki TAKAHASHI and Toshiaki TSUNOGAE